

Temporal trends and demographic patterns in hypertension-related mortality with intracerebral hemorrhage in the United States: 1999–2020

Paraag Vashist, Khansa Younus, Sindiri Rohit, Sowkarthick K S, Abhishek Hanumanpratap Singh Kshatri, Surya Vamsi Gadde

DOI: <https://doi.org/10.21542/gcsp.2026.3>
VOLUME: 2026
ISSUE: 1
ARTICLE: 3

PMC PUB. DATE: 28 February 2026
PMC COLLECTION DATE: 28 February 2026
PMC ARTICLE TYPE: research-article

ARTICLE TYPE: Research article
SUBMITTED: 18 Oct 2025
ACCEPTED: 22 Jan 2026
RUNNING HEAD: Vashist et al., GCSP 2026:3

CITE THIS ARTICLE AS: Vashist P, Younus K, Rohit S, Sowkarthick KS, Kshatri AHS, Gadde SV. Temporal trends and demographic patterns in hypertension-related mortality with intracerebral hemorrhage in the United States: 1999–2020, Global Cardiology Science and Practice 2026:3 <https://doi.org/10.21542/gcsp.2026.3>

COPYRIGHT: 2026 The Author(s), licensee Magdi Yacoub Institute. This is an open access article distributed under the terms of the Creative Commons Attribution license CC BY-4.0, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Please note this is an 'Article in Press' and will be replaced with a typeset PDF & PubMed Central deposit in the coming weeks. Citation details and DOI will remain the same.

Temporal trends and demographic patterns in hypertension-related mortality with intracerebral hemorrhage in the United States: 1999–2020

Paraag Vashist¹, Khansa Younus², Sindiri Rohit³, Sowkarthick K S^{4,*}, Abhishek Hanumanpratap Singh Kshatri⁵, Surya Vamsi Gadde⁶

1. *Department of General Internal Medicine, Leighton Hospital, Mid Cheshire Hospital, NHS Foundation Trust*
2. *Department of Family Medicine, Carmichael Medical Clinic.*
3. *Department of Internal Medicine, All India Institute of Medical Sciences*
4. *Department of Critical Care, St John's Medical College Hospital, Bangalore*
5. *Department of Emergency Medicine, Apollo Hospitals*
6. *Department of Internal Medicine, Gayatri Vidya Parishad Institute of Health care and Medical Technology.*

*sowkarthicksaravanan@gmail.com

Abstract

Introduction: Hypertension contributes to cardiovascular mortality, yet its role as an underlying cause with intracerebral hemorrhage (ICH) as a contributing cause is underexplored. Understanding demographic and geographic patterns can guide prevention.

Aim: To analyze mortality trends and demographic disparities in deaths with hypertension as the underlying cause and ICH as a contributing cause, using the CDC Multiple Cause of Death (MCD) database from 1999–2020.

Methods: Retrospective observational analysis of the CDC MCD database for adults aged ≥ 25 years in the United States from 1999–2020. Deaths were included if hypertension (I10–I15) was the underlying cause and intracerebral hemorrhage (ICH I61) a contributing cause. Data were stratified by gender, race, geographic region, and place of death. Crude and age-adjusted mortality rates per 1,000,000 and annual percentage change (APC) were calculated. Temporal trends were assessed using Joinpoint software.

Results: Among 14,613 deaths (crude rate 3.3 per million), most decedents were male (56.5%)

and White (62.8%); Black/African American individuals accounted for 31.2%, Asian/Pacific Islander 5.3%, and American Indian/Alaska Native 0.7%. Most deaths occurred in metropolitan areas (89.4%) and in medical facilities (61.0%), while 25.2% occurred at home. Age-adjusted rates were stable from 1999–2007, declined 2007–2013, then rose 2013–2020. Females showed larger early declines followed by a modest rise, whereas males experienced a later sharper increase. Black/African American decedents had early declines with a subsequent plateau, while White decedents showed smaller early change, followed by decline and later increase.

Conclusion: Deaths with hypertension as the underlying cause and ICH as a contributing cause were concentrated among males, White and Black populations, and metropolitan residents, with most deaths in medical facilities. Notably, after a period of decline, mortality rates increased again after 2013, particularly among males and White individuals, underscoring the urgency of renewed prevention efforts.

Introduction

Hypertension refers to the force exerted by circulating blood against the arterial walls, defined as systolic blood pressure ≥ 140 mmHg and/or diastolic blood pressure ≥ 90 mmHg [1]. Treatment across all age groups significantly reduces cardiovascular and cerebrovascular risk [1]. In the U.S. population, the prevalence of hypertension among adults aged over 20 years is 48.4% [2]. The age-adjusted mortality rate increased from 2.8 to 9.4 per 100,000 from 1999 to 2019, reaching 13.9 per 100,000 by 2021, with younger adults experiencing particularly sharp increases in recent years [3]. Globally, hypertension contributes to an estimated 10.4 million deaths annually [4].

Modifiable risk factors for hypertension include excess body weight, physical inactivity, alcohol use, smoking, diabetes mellitus, obstructive sleep apnea, and chronic kidney disease, while non-modifiable risk factors consist of age, race/ethnicity, and family history [5].

Intracerebral hemorrhage (ICH) is a type of stroke characterized by bleeding within the brain parenchyma, with or without ventricular extension. Chronic hypertension is the most important risk factor, accounting for over 60% of ICH cases [6]. Hypertensive ICH typically involves deep brain structures, such as the basal ganglia, thalamus, pons, and cerebellum, due to small-vessel pathology, including lipohyalinosis and microaneurysm rupture [7]. Of all strokes in the United States, ICH accounts for approximately 10–20% and represents a particularly devastating subtype, with a 30-day mortality rate approaching 40% and a high rate of long-term disability among survivors [8].

In the United States, the combined mortality burden of hypertension and ICH remains substantial; however, most prior research has examined these conditions in isolation [9,10]. While ICH is commonly recorded as the underlying cause of death in hemorrhagic stroke

cases, death certification practices also recognize clinical scenarios in which long-standing, poorly controlled hypertension is identified as the initiating condition in the causal chain leading to death, with ICH documented as a downstream or contributing event. Such scenarios may include patients with chronic hypertensive vasculopathy who experience fatal intracerebral hemorrhage, where certifying physicians attribute the death primarily to hypertension as the root etiologic driver rather than to the acute hemorrhagic event alone.

Accordingly, in this study, we examine mortality in which hypertension (ICD-10 codes I10–I15) is recorded as the underlying cause of death, and intracerebral hemorrhage (ICD-10 code I61) is listed as a contributing cause of death. This analytic approach provides a hypertension-centric lens that emphasizes deaths attributable to chronic hypertensive disease processes while capturing ICH as a clinically significant and often fatal complication. Although less commonly examined than stroke-centric analyses in which ICH is the underlying cause, this perspective is particularly appropriate for evaluating hypertension-related mortality, as it highlights the role of long-term blood pressure control failure in precipitating fatal cerebrovascular outcomes and allows assessment of temporal and demographic patterns in deaths explicitly attributed to hypertension at the population level.

Aim and objectives

To evaluate U.S. mortality where hypertensive diseases (I10–I15) were the underlying cause and intracerebral haemorrhage ICH (I61) a contributing cause among adult's ≥ 25 years (1999–2020), reporting totals, crude rate per 1,000,000, and distributions by sex, race, 2013 NCHS urban–rural category, and place of death using CDC WONDER MCD.

Methodology

Data were extracted from the CDC-WONDER Multiple Cause of Death (MCD) database for retrospective analysis [11]. The process of data extraction was performed on 3rd August 2025. The database encompasses death certificate records spanning 1999 to 2020 and was queried to include individuals aged 25 years and older, given the well-documented rarity of hypertension-associated intracranial hemorrhage below this age threshold. As the CDC-WONDER MCD database contains fully de-identified, publicly available data, this study was exempt from formal ethics committee review and institutional approval [12].

Hypertension (I10–I15) was designated as the underlying cause of death, with intracranial hemorrhage (I61) specified as a contributing multiple cause of death.

Demographic variables included sex (male and female) and race/ethnicity (American Indian or

Alaska Native, Asian or Pacific Islander, Black or African American, and White), reflecting the predominant racial groups represented within the United States population. These variables were incorporated to evaluate potential disparities in mortality outcomes across demographic subgroups.

Geographic variables were defined according to the 2013 National Center for Health Statistics (NCHS) urbanization classification scheme, which stratifies counties into six categories: large central metro, large fringe metro, medium metro, and small metro within the metropolitan designation, and micropolitan and non-core areas within the non-metropolitan designation[13].

Place of death was further classified into five predefined categories: medical facility (inpatient), medical facility (outpatient), decedent's home, hospice facility, and nursing home [14]. Age-standardized mortality rates were calculated per 1,000,000 population using the United States 2000 standard population as the reference, facilitating valid and accurate temporal comparisons across the study period [15].

Descriptive statistics were generated for each variable as absolute frequencies and proportions using CDC-WONDER MCD data in accordance with the specified extraction criteria; variables failing to meet these criteria were excluded from the analysis. Temporal trends in age-adjusted mortality rates from 1999 to 2020 were subsequently examined using Joinpoint regression software, which enabled the calculation of annual percentage changes and the identification of statistically significant inflection points over the study period [16].

Results

Between 1999 and 2020, the CDC Multiple Causes of Death (MCD) database documented 14,613 fatalities in the United States among individuals aged 25 years and older attributable to hypertension (ICD-10: I10–I15). This analysis also incorporated cases where intracerebral hemorrhage (ICD-10: I61) was recorded as a multiple cause of death, in order to account for cerebrovascular complications associated with hypertension. The crude mortality rate for hypertension was found to be 3.3 per 1,000,000 populations. Deaths not meeting these criteria were excluded from the analysis.

Demographic and geographical characteristics

Among 14,613 individuals included in the mortality dataset between 1999 and 2020 with hypertension listed as the underlying cause of death and intracerebral hemorrhage as a contributing cause, males represented a greater proportion of deaths than females (8,258 [56.5%] vs. 6,355 [43.5%]), reflecting the consistently higher hypertension and hemorrhagic stroke burden reported among men in epidemiologic studies.

The age distribution demonstrated that mortality clustered predominantly in mid-to-late adulthood, with the highest proportions observed in the 45–54 years (3,262; 22.3%) and 55–64 years (3,100; 21.2%) groups, followed by those aged ≥ 75 years. Fewer deaths occurred among adults younger than 45 years (total 1,623; 11.1%), indicating that hypertension-related intracerebral hemorrhage remains largely a condition affecting older age strata.

Racial patterns showed that White individuals accounted for the majority of deaths (9,183; 62.8%); however, Black or African American individuals comprised nearly one-third of deaths (4,561; 31.2%), a proportion notably higher than their representation in the U.S. population, suggesting an unequal burden likely driven by longstanding disparities in hypertension control and access to preventive care.

Most deaths occurred in metropolitan settings (13,065; 89.4%), consistent with population clustering in these regions, but potentially also reflective of greater diagnostic capture and access to acute care services where hemorrhagic stroke is more readily identified. Taken together, these demographic patterns illustrate a concentration of mortality among older adults, males, and Black populations, underscoring the need for targeted risk-reduction efforts in high-burden demographic groups (Table 1).

Mortality rate comparisons

Age-adjusted mortality rates showed substantial demographic variation (Table 2). Overall, mortality was higher among males than females (3.8 vs 2.4 per 100,000), reflecting a greater burden of hypertension-associated ICH deaths in men. Black/African American adults experienced the highest mortality (8.7 per 100,000), nearly four-fold higher than White adults (2.3 per 100,000), indicating persistent racial disparities. Asian/Pacific Islander adults had intermediate mortality (3.7 per 100,000), while American Indian/Alaska Native adults demonstrated lower but imprecise estimates due to small counts (2.3 per 100,000; 95% CI 1.9–2.9).

Mortality was modestly higher in metropolitan than non-metropolitan areas (2.1 vs 1.3 per 100,000), a pattern consistent with the concentration of stroke care and hypertension burden in urban regions. These findings highlight demographic inequalities in hypertension-related ICH mortality and justify subgroup trend analyses.

Temporal trends

From 1999–2020, overall age-adjusted mortality due to hypertension with intracerebral hemorrhage as a contributing cause demonstrated three statistically distinct trend segments (Figure 1, Table 3). Mortality rates remained stable between 1999–2007 (APC = 0.05%; 95% CI –1.09 to 3.28; $p = 0.86$), followed by a significant decline from 2007–2013 (APC = –4.31%; 95% CI –9.41 to –2.30; $p = 0.006$). However, this reduction reversed thereafter, with rates increasing significantly from 2013–2020 (APC = 2.66%; 95% CI 0.94 to 7.05; $p = 0.008$), indicating a resurgence in hypertension-related mortality after an earlier period of improvement.

Sex-stratified analyses revealed divergent patterns between males and females (Figure 2). In females, mortality trends were initially stable from 1999–2007 (APC = -0.57% ; 95% CI -2.35 to 6.22 ; $p = 0.73$), followed by a significant decline from 2007–2012 (APC = -6.70% ; 95% CI -12.66 to -3.08 ; $p = 0.02$), and subsequently a significant rise during 2012–2020 (APC = 2.32% ; 95% CI 0.37 to 7.25 ; $p = 0.03$). In contrast, males experienced stable rates from 1999–2007 (APC = 0.47% ; 95% CI -0.65 to 4.77 ; $p = 0.42$), a significant decline between 2007–2015 (APC = -2.82% ; 95% CI -8.35 to -1.52 ; $p = 0.005$), and a subsequent sharper increase from 2015–2020 (APC = 4.96% ; 95% CI 1.98 to 12.50 ; $p = 0.004$). These contrasting trajectories were statistically non-parallel ($p = 0.010$), confirming sex-specific differences in temporal mortality patterns (Table 4).

Race-stratified analyses further demonstrated heterogeneity in mortality trends (Figure 3). Among Black/African American adults, mortality declined during 1999–2008 (APC = -2.15% ; 95% CI -3.54 to 2.56 ; $p = 0.13$) and more sharply between 2008–2014 (APC = -7.23% ; 95% CI -13.58 to -4.66 ; $p = 0.016$), before stabilizing during 2014–2020 (APC = 0.22% ; 95% CI -2.66 to 9.88 ; $p = 0.74$). Among White adults, mortality was stable from 1999–2008 (APC = 0.05% ; 95% CI -1.43 to 1.55 ; $p = 0.94$), followed by a non-significant decline during 2008–2013 (APC = -3.00% ; 95% CI -7.89 to 2.14 ; $p = 0.23$), and a significant rise between 2013–2020 (APC = 3.34% ; 95% CI 1.11 to 5.62 ; $p = 0.006$). Comparable trend testing confirmed significant non-parallelism between Black and White groups ($p = 0.0002$; Table 4), indicating persistent racial disparities and differing temporal dynamics. Trends for American Indian/Alaska Native and Asian/Pacific Islander adults could not be assessed due to data suppression for small counts (<10).

Urban–rural patterns also differed significantly over time ($p = 0.0036$; Table 4), with metropolitan areas showing a significant decline from 2008–2013 (APC = -5.41% ; 95% CI -9.40 to -3.07 ; $p < 0.001$) followed by an increase from 2013–2020 (APC = 2.05% ; 95% CI 0.66 to 4.54 ; $p = 0.004$). In contrast, non-metropolitan trends were more variable, with an initial decline followed by a short-term rise and subsequent non-significant decrease; however, suppression of unstable estimates limited interpretation. Taken together, these results highlight a nationwide rebound in hypertension-related intracerebral hemorrhage mortality after 2013, with significant differences by sex, race, and urbanicity.

In a complementary analysis examining deaths in which intracerebral hemorrhage was recorded as the underlying cause of death and hypertension as a contributing cause, age-adjusted mortality trends differed from those observed in the primary analysis (Figure 4). Mortality rates increased sharply between 1999 and 2001 (APC = $+458.64\%$; 95% CI not shown; $p < 0.05$), followed by a significant decline from 2001 to 2004 (APC = -16.80% ; $p < 0.05$). From 2004 through 2020, a continued but more gradual decline in age-adjusted mortality was observed (APC = -1.74% ; $p < 0.05$). No post-2013 increase in mortality rates was identified in this stroke-centric analysis, indicating a sustained downward trend when

intracerebral hemorrhage was specified as the underlying cause of death and hypertension was listed as a contributing condition.

Discussion

This population-based study demonstrates important long-term trends in mortality attributed to hypertension (ICD-10 codes I10–I15) with intracerebral hemorrhage (ICH; ICD-10 code I61) as a contributing cause of death among United States adults aged 25 years and older over the two-decade period from 1999 to 2020.

Over these 22 years, 14,613 deaths were identified, with a distinct three-phase temporal pattern: an initial period of stability, a significant decline between 2007 and 2013, followed by a statistically significant increase in mortality after 2013. This post-2013 reversal represents the most clinically and public health–relevant finding of the study.

Mortality reversal after 2013

The resurgence in age-adjusted mortality observed after 2013 likely reflects a convergence of biological, clinical, and healthcare-system factors. Earlier declines in mortality may be attributable to improved hypertension awareness, treatment, and acute stroke care, consistent with prior national trends [17,18]. These favorable trends also coincided with broader healthcare system reforms that expanded insurance coverage and access to preventive and chronic disease care, which have been associated with improvements in population-level health outcomes and reductions in mortality over time [19].

However, sustained long-term blood pressure control remains challenging, particularly in populations with high cardiometabolic risk burdens and adverse social determinants of health. Even with improved access to care, gaps in hypertension treatment adherence, continuity of care, and effective blood pressure control persist, especially among socioeconomically disadvantaged groups [20]. Chronic poorly controlled hypertension leads to maladaptive shifts in cerebral autoregulation, structural weakening of small penetrating arteries, and development of Charcot–Bouchard microaneurysms, all of which increase susceptibility to intracerebral hemorrhage [21–23].

The observed post-2013 reversal may also coincide with broader changes in hypertension management paradigms, evolving clinical thresholds for treatment intensity, and the increasing prevalence of obesity, diabetes, and chronic kidney disease—factors known to amplify long-term cerebrovascular risk [24, 25]. In parallel, the expanding use of anticoagulant therapies in aging hypertensive populations may increase the severity and fatality of hemorrhagic events when blood pressure control is suboptimal [24,26,27]. Although these therapies reduce thromboembolic risk, their interaction with poorly controlled hypertension may disproportionately worsen outcomes once intracerebral hemorrhage occurs.

Importantly, the complementary analysis in which ICH was treated as the underlying cause demonstrated a continued long-term decline in stroke-centric mortality, suggesting that the observed post-2013 increase is specific to deaths attributed primarily to hypertension rather than hemorrhagic stroke overall. This divergence underscores the importance of cause-of-death attribution and supports prior evidence showing that insurance expansion and improved access to cardiovascular care are associated with reductions in overall cardiac and cerebrovascular mortality outside acute care settings, even when condition-specific or attribution-based mortality trends differ [28].

Racial disparities

Marked racial disparities were evident throughout the study period. While White individuals accounted for the highest absolute number of deaths, Black/African American adults experienced substantially higher age-adjusted mortality rates. This finding is consistent with prior literature demonstrating an earlier onset of hypertension, greater disease severity, and poorer blood pressure control among Black populations [29,30]. Importantly, these disparities cannot be attributed solely to biological susceptibility and instead reflect the cumulative effects of adverse social determinants of health, including socioeconomic disadvantage, chronic psychosocial stress, and neighborhood-level inequities.

Structural racism plays a critical role in shaping long-term cardiovascular risk through mechanisms such as residential segregation, reduced access to healthy food environments, limited opportunities for physical activity, and sustained exposure to stressors that contribute to poor hypertension control. Differences in healthcare access, continuity of care, and quality of hypertension management—including delayed diagnosis, undertreatment, and reduced access to specialty stroke care—may further increase vulnerability to catastrophic cerebrovascular outcomes among Black adults. Prior population-based studies have demonstrated that Black individuals experience a disproportionately higher risk of intracerebral hemorrhage and stroke-related mortality, even after accounting for traditional risk factors [31,32].

Our findings align with a broader body of evidence documenting persistent racial inequities in hypertension prevalence, treatment, and cardiovascular mortality in the United States [26–28]. Interpretation of trends among American Indian/Alaska Native and Asian/Pacific Islander populations was limited by data suppression due to small counts, underscoring ongoing gaps in national surveillance systems for smaller racial and ethnic groups and highlighting the need for improved data capture to better understand disparities in these populations.

Urban–rural differences

Geographic disparities were also evident, with the majority of deaths occurring in metropolitan areas. While urban populations may have a higher burden of cardiometabolic risk factors and social stressors, differences in mortality patterns between metropolitan and non-metropolitan areas likely extend beyond lifestyle factors alone. Access to timely emergency medical services, hospital availability, and the distribution of comprehensive

stroke centers vary substantially by geography and may critically influence outcomes following intracerebral hemorrhage [33,34].

Non-metropolitan populations often face longer emergency medical services response and transport times, fewer hospitals with advanced stroke capabilities, and limited access to specialized neurocritical care, all of which may adversely affect survival following acute cerebrovascular events [35]. Conversely, higher mortality observed in metropolitan areas may reflect greater disease burden, population density, and differential healthcare utilization patterns, including higher rates of diagnostic recognition and cause-of-death attribution.

-

Additionally, regional variation in death certificate completion practices may influence whether hypertension and intracerebral hemorrhage are recorded as underlying versus contributing causes of death, potentially contributing to observed geographic differences [36]. Together, these findings suggest that both urban and rural populations require tailored, context-specific strategies addressing healthcare infrastructure, equitable access to emergency stroke care, and long-term hypertension prevention and management.

Limitations

This study has limitations inherent to analyses of death certificate–based data. Cause-of-death reporting may be subject to misclassification bias due to variability in physician certification practices and inaccuracies in documenting the causal sequence of death. Although ICD-10 codes were applied consistently, changes in diagnostic awareness, documentation, and coding practices over time may have influenced observed trends. As a population-level analysis, findings are susceptible to ecological fallacy and cannot be interpreted as individual-level causal associations. The absence of individual clinical information—including blood pressure control, medication adherence, comorbidities, anticoagulant use, and access to acute care—limits adjustment for confounders. Small cell counts and data suppression (<10 deaths) restricted reliable estimation for certain racial groups and limited calculation of subgroup-specific trend and comparative statistics. Declining autopsy rates may have reduced diagnostic certainty, and COVID-19–related disruptions in 2020 may have affected mortality reporting and completeness.

Conclusions

From 1999 to 2020, 14,613 deaths occurred in the United States in which hypertensive disease was recorded as the underlying cause of death and intracerebral hemorrhage as a contributing cause, corresponding to a crude mortality rate of 3.3 per 1,000,000 adults. Mortality was disproportionately concentrated among men, metropolitan residents, and White and Black populations, with most deaths occurring in inpatient medical facilities. After a period of decline

between 2007 and 2013, age-adjusted mortality rates increased significantly thereafter, indicating a reversal of earlier gains in hypertension-related mortality reduction.

This resurgence in mortality is clinically and public-health relevant, as it suggests gaps in long-term blood pressure control, risk-factor modification, and continuity of care despite advances in antihypertensive therapy and stroke management. Clinically, these findings underscore the need for earlier identification of high-risk individuals, sustained hypertension control across the life course, and aggressive management of comorbid conditions that increase susceptibility to intracerebral hemorrhage. From a public health perspective, targeted prevention strategies—including population-level blood pressure screening, improved adherence to hypertension treatment guidelines, and focused interventions in high-burden demographic and metropolitan populations—are essential to prevent further increases in mortality.

Future efforts should prioritize strengthening primary prevention, improving equitable access to chronic cardiovascular care, and monitoring hypertension-related outcomes using cause-of-death frameworks that capture both underlying disease and fatal cerebrovascular complications. Addressing the observed resurgence is critical to reversing current trends and reducing preventable deaths attributable to hypertension and intracerebral hemorrhage.

References

1. Ramzy, I. (2019). Definition of hypertension and pressure goals during treatment (ESC-ESH Guidelines 2018). *E-Journal of Cardiology Practice*, 17(18). European Society of Cardiology.
2. Chobufo, M. D., Gayam, V., Solunoy, J., Rahman, E. U., Enoru, S., Foryoung, J. B., Agbor, V. N., Dufresne, A., & Nfor, T. (2020). Prevalence and control rates of hypertension in the USA: 2017–2018. *International Journal of Cardiology Hypertension*, 6, 100044. <https://doi.org/10.1016/j.ijchy.2020.100044>
3. Siddiqi AK, Ali KM, Maniya MT, Rashid AM, Khatri SA, Garcia M, Quintana RA, NaeemM. The hidden epidemic: Hypertension-related mortality surges amongst younger adults in the United States. *Curr Probl Cardiol*. 2024 Dec;49(12):102842. doi: 10.1016/j.cpcardiol.2024.102842. Epub 2024 Sep 12. PMID: 39270766.
4. Nabi, R., Zanub, A., Akhtar, M., et al. (2025). Concomitant mortality trends due to obesity and hypertension in the U.S.: A 20-year retrospective analysis of the CDC WONDER database. *BMC Cardiovascular Disorders*, 25, 496. <https://doi.org/10.1186/s12872-025-04909-z>
5. Whelton PK, Carey RM, Aronow WS, Casey DE Jr, Collins KJ, Dennison Himmelfarb C, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: A report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018 Jun;71(6):e13–115. PMID:29133356
6. Rajashekar D, Liang JW. Intracerebral Hemorrhage. [Updated 2023 Feb 6]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan-. Available from:

- <https://www.ncbi.nlm.nih.gov/books/NBK553103/>
7. Auer, R. N., & Sutherland, G. R. (2006). Primary intracerebral hemorrhage: Pathophysiology. *The Canadian Journal of Neurological Sciences*, 32(Suppl 2), S3–S12.
<https://doi.org/10.1017/S0317167100004161>
 8. Morotti, A., & Goldstein, J. N. (2016). Diagnosis and management of acute intracerebral hemorrhage. *Emergency Medicine Clinics of North America*, 34(4), 883–899.
<https://doi.org/10.1016/j.emc.2016.06.010>
 9. Ford, E. S. (2011). Trends in mortality from all causes and cardiovascular disease among hypertensive and nonhypertensive adults in the United States. *Circulation*, 123(16), 1737–1744. <https://doi.org/10.1161/CIRCULATIONAHA.110.005645>
 10. DeLago, A. J., Jr., Singh, H., Jani, C., Rupal, A., Shalhoub, J., Goodall, R., Shah, D., Hartley, A., Hammond-Haley, M., Marshall, D. C., Gizzi, M., AbdelRazek, M., & Saliccioli, J. D. (2022). An observational epidemiological study to analyze intracerebral hemorrhage across the United States: Incidence and mortality trends from 1990 to 2017. *Journal of Stroke and Cerebrovascular Diseases*, 31(4), 106216. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2021.106216>
 11. Centers for Disease Control and Prevention, National Center for Health Statistics. National Vital Statistics System, Mortality 1999–2020 on CDC WONDER Online Database, released in 2021. Data are from the Multiple Cause of Death Files, 1999–2020, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Accessed at <http://wonder.cdc.gov/mcd-icd10.html> on Aug 15, 2025 3:33:30 PM.
 12. DeLusignan S, Liyanage H, Di Iorio CT, Chan T, Liaw ST. Using routinely collected health data for surveillance, quality improvement and research: Framework and key questions to assess ethics and privacy and enable data access. *BMJ Health & Care Informatics*. 2015 Oct 1;22(4).
 13. Ingram, D. D., & Franco, S. J. (2014, April). 2013 NCHS Urban–Rural Classification Scheme for Counties. *Vital and Health Statistics: Series 2, Number 166*. National Center for Health Statistics.
 14. National Center for Health Statistics. (n.d.). Underlying cause of death, 1999–2020: Data definitions – Place of death. Centers for Disease Control and Prevention.
<https://wonder.cdc.gov/wonder/help/ucd.html>
 15. Sorlie PD, Thom TJ, Manolio T, et al: Age-adjusted death rates: consequences of the Year 2000 standard. *Ann Epidemiol*. 1999, 9:93-100. 10.1016/s1047-2797(98)00061-1
 16. Joinpoint Regression Program, Version 5.0.2 - May 2023; Statistical Methodology and Applications Branch, Surveillance Research Program, National Cancer Institute. Accessed October 29, 2023. <https://surveillance.cancer.gov/joinpoint/>.
 17. Noushad MA, Mulholland E, Jaisukhalal J, Abukhater R, Qayyum W, Mukesh MA, Jerome D Silva J, Sheikh MT, Arshad MM, Essa NE, Zulfiqar E, Ahmed M, Ahmed R. Trends and Disparities in Hemorrhagic Stroke and Hypertension-Related Mortality in the United States From 1999 to 2023: A CDC WONDER Database Analysis. *Brain Behav*. 2025 Jul;15(7):e70704. doi: 10.1002/brb3.70704. PMID: 40686020; PMCID: PMC12277664.
 18. Saha, R., Al-Salihi, M.M., Elazim, A.A., Solanki, D., Kapuria, P. and Dalal, S.S. (2025), Trends in Intracranial Hemorrhage Mortality in the U.S.: A 22-Year Analysis (1999–2020). *Neurol Clin Neurosci*, 13: 174-180. <https://doi.org/10.1111/ncn3.70000>
 19. Aparna Soni, Laura R. Wherry, and Kosali I. Simon. How Have ACA Insurance Expansions Affected Health Outcomes? Findings From The Literature. *Health Affairs* 2020 39:3, 371-378. DOI: Aparna Soni, Laura R. Wherry, and Kosali I. Simon

20. H A, Bb G, K F, et al. Role of health insurance and neighborhood-level social deprivation on hypertension control following the affordable care act health insurance opportunities. *Soc Sci Med.* 2020;265:113439. doi:10.1016/j.socscimed.2020.113439
21. Silverman A, Petersen NH. Physiology, Cerebral Autoregulation. 2023 Mar 15. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan–. PMID: 31985976.
22. Dandapani BK, Suzuki S, Kelley RE, Reyes-Iglesias Y, Duncan RC. Relation between blood pressure and outcome in intracerebral hemorrhage. *Stroke.* 1995 Jan;26(1):21-4. doi: 10.1161/01.str.26.1.21. PMID: 7839391.
23. Gupta K, Das JM. Charcot-Bouchard Aneurysm. 2023 Jul 17. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan–. PMID: 31971704.
24. Ikram MA, Wieberdink RG, Koudstaal PJ. International epidemiology of intracerebral hemorrhage. *Curr Atheroscler Rep.* 2012 Aug;14(4):300-6. doi: 10.1007/s11883-012-0252-1. PMID: 22538431; PMCID: PMC3388250.
25. Biffi A, Cortellini L, Nearnberg CM, Ayres AM, Schwab K, Gilson AJ, Rost NS, Goldstein JN, Viswanathan A, Greenberg SM, Rosand J. Body mass index and etiology of intracerebral hemorrhage. *Stroke.* 2011 Sep;42(9):2526-30. doi: 10.1161/STROKEAHA.111.617225. Epub 2011 Jul 21. PMID: 21778442; PMCID: PMC3164290.
26. O'Donnell MJ, Xavier D, Liu L, Zhang H, Chin SL, Rao-Melacini P, Rangarajan S, Islam S, Pais P, McQueen MJ, Mondo C, Damasceno A, Lopez-Jaramillo P, Hankey GJ, Dans AL, Yusuf K, Truelsen T, Diener HC, Sacco RL, Ryglewicz D, Czlonkowska A, Weimar C, Wang X, Yusuf S; INTERSTROKE investigators. Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): a case- control study. *Lancet.* 2010 Jul 10;376(9735):112-23. doi: 10.1016/S0140- 6736(10)60834-3. Epub 2010 Jun 17. PMID: 20561675.
27. Ariesen MJ, Claus SP, Rinkel GJ, Algra A. Risk factors for intracerebral hemorrhage in the general population: a systematic review. *Stroke.* 2003 Aug;34(8):2060-5. doi: 10.1161/01.STR.0000080678.09344.8D. Epub 2003 Jul 3. PMID: 12843354.
28. Ginger Y. Jiang, MD, MBA, John W. Urwin, MD, and Jason H. Wasfy, MD, MPhil. Medicaid Expansion Under the Affordable Care Act and Association With Cardiac Care: A Systematic Review. *Circulation: Cardiovascular Quality and Outcomes.* Volume 16, Number 6 <https://doi.org/10.1161/CIRCOUTCOMES.122.009753>
29. Poorthuis MH, Algra AM, Algra A, Kappelle LJ, Klijn CJ. Female- and Male- Specific Risk Factors for Stroke: A Systematic Review and Meta-analysis. *JAMA Neurol.* 2017 Jan 1;74(1):75-81. doi: 10.1001/jamaneurol.2016.3482. PMID: 27842176.
30. Aggarwal R, Chiu N, Wadhwa RK, Moran AE, Raber I, Shen C, Yeh RW, Kazi DS. Racial/Ethnic Disparities in Hypertension Prevalence, Awareness, Treatment, and Control in the United States, 2013 to 2018. *Hypertension.* 2021 Dec;78(6):1719-1726. doi: 10.1161/HYPERTENSIONAHA.121.17570. Epub 2021 Aug 9. PMID: 34365809; PMCID: PMC10861176.
31. Alruwaili A, Alanazy ARM. Prehospital Time Interval for Urban and Rural Emergency Medical Services: A Systematic Literature Review. *Healthcare (Basel).* 2022;10(12):2391. Published 2022 Nov 29. doi:10.3390/healthcare10122391
32. Kittner SJ, Sekar P, Comeau ME, et al. Ethnic and Racial Variation in Intracerebral Hemorrhage Risk Factors and Risk Factor Burden. *JAMA Netw Open.* 2021;4(8):e2121921. Published 2021 Aug

2. doi:10.1001/jamanetworkopen.2021.21921
33. Whelton PK, Einhorn PT, Muntner P, Appel LJ, Cushman WC, Diez Roux AV, Ferdinand KC, Rahman M, Taylor HA, Ard J, Arnett DK, Carter BL, Davis BR, Freedman BI, Cooper LA, Cooper R, Desvigne-Nickens P, Gavini N, Go AS, Hyman DJ, Kimmel PL, Margolis KL, Miller ER 3rd, Mills KT, Mensah GA, Navar AM, Ogedegbe G, Rakotz MK, Thomas G, Tobin JN, Wright JT, Yoon SS, Cutler JA; National Heart, Lung, and Blood Institute Working Group on Research Needs to Improve Hypertension Treatment and Control in African Americans. Research Needs to Improve Hypertension Treatment and Control in African Americans. *Hypertension*. 2016 Nov;68(5):1066-1072. doi: 10.1161/HYPERTENSIONAHA.116.07905. Epub 2016 Sep 12. PMID:27620388; PMCID: PMC5063700.
 34. Kamin Mukaz D, Dawson E, Howard VJ, Cushman M, Higginbotham JC, Judd SE, Kissela BM, Safford MM, Soliman EZ, Howard G. Rural/urban differences in the prevalence of stroke risk factors: A cross-sectional analysis from the REGARDS study. *J Rural Health*. 2022 Jun;38(3):668-673. doi: 10.1111/jrh.12608. Epub 2021 Jul 16. PMID: 34270125; PMCID: PMC8761227.
 35. Qureshi AI, Giles WH, Croft JB. Racial differences in the incidence of intracerebral hemorrhage: effects of blood pressure and education. *Neurology*. 1999;52(8):1617-1621. doi:10.1212/wnl.52.8.1617
 36. Hammond G, Luke AA, Elson L, Towfighi A, Joynt Maddox KE. Urban-Rural Inequities in Acute Stroke Care and In-Hospital Mortality. *Stroke*. 2020;51(7):2131-2138. doi:10.1161/STROKEAHA.120.029318

TABLES:

Table 1. Baseline demographic characteristics of decedents with hypertension as the underlying cause of death and intracerebral hemorrhage as a contributing cause, United States, 1999–2020.

Baseline demographic distribution of decedents aged ≥25 years in whom hypertension (ICD-10 I10–I15) was recorded as the underlying cause of death and intracerebral hemorrhage (ICD-10 I61) as a contributing cause, using CDC WONDER Multiple Cause of Death data (1999–2020). Percentages are calculated within each category.

Characteristic	No. of deaths	Percentage (%)
Total	14613	100
Sex		
Male	8258	56.50
Female	6355	43.50
Age, in years		
25–34	263	1.80
35 – 44	1360	9.30
45 – 54	3262	22.30
55 – 64	3100	21.20
65–74	2063	14.10
75 – 84	2516	17.20
≥85	2049	14.00
Race		
American Indian or Alaska Native	96	0.70
Asian or Pacific Islander	773	5.30
Black or African American	4561	31.20
White	9183	62.80
Residence		
Metropolitan	13065	89.40
Non-metropolitan	1548	10.60

Table 2. Age-adjusted mortality rates by demographic characteristics for hypertension-related intracerebral hemorrhage mortality, United States, 1999–2020.

Age-adjusted mortality rates (AAMR) per 100,000 population were calculated using the direct method and standardized to the 2000 U.S. standard population. Rates represent deaths with hypertension as the underlying cause of death and intracerebral hemorrhage as a contributing cause, derived from CDC WONDER (1999–2020).

Demographic characteristic	AAMR (per 1,000,000)	95% Confidence Interval
Sex		
Male	3.8	3.8 - 3.9
Female	2.4	2.4 - 2.5
Race		

White	2.3	2.3 - 2.3
Black/African American	8.7	8.5 - 9
American Indian/Alaska Native	2.3	1.9 - 2.9
Asian/Pacific Islander	3.7	3.4 - 4
Residence		
Metropolitan	2.1	2.1 - 2.2
Non-metropolitan	1.3	1.2 - 1.3

Table 3. Joinpoint regression analysis of age-adjusted mortality trends for hypertension-related intracerebral hemorrhage mortality, United States, 1999–2020. * Indicates that the APC (Annual percentage change) is significantly differ from zero at alpha=0.05 level.

Cohort	Segment	Lower Endpoint	Upper Endpoint	APC	95% CI	P-Value
Overall	1	1999	2007	0.0512	-1.09 - 3.28	0.8582
Overall	2	2007	2013	-4.3115*	-9.41 - 2.3	0.0064
Overall	3	2013	2020	2.6573*	0.94 - 7.05	0.0080
Female - 2 Joinpoints	1	1999	2007	-0.5656	-2.35 - 6.22	0.7267
Female - 2 Joinpoints	2	2007	2012	-6.6956*	-12.66 - 3.08	0.0200
Female - 2 Joinpoints	3	2012	2020	2.3221*	0.37 - 7.25	0.0324
Male - 2 Joinpoints	1	1999	2007	0.4677	-0.65 - 4.77	0.4183
Male - 2 Joinpoints	2	2007	2015	-2.8157*	-8.35 - 1.52	0.0052
Male - 2 Joinpoints	3	2015	2020	4.9570*	1.98 - 12.5	0.0036
Black or African American - 2 Joinpoints	1	1999	2008	-2.1455	-3.54 - 2.56	0.1252
Black or African American - 2 Joinpoints	2	2008	2014	-7.2279*	-13.58 - 4.66	0.0164
Black or African American - 2 Joinpoints	3	2014	2020	0.2174	-2.66 - 9.88	0.7351
White - 2 Joinpoints	1	1999	2008	0.0499	-1.43 - 1.55	0.9438
White - 2 Joinpoints	2	2008	2013	-3.0030	-7.89 - 2.14	0.2262
White - 2 Joinpoints	3	2013	2020	3.3401*	1.11 - 5.62	0.0061
Metro - 2 Joinpoints	1	1999	2008	0.1990	-0.73 - 1.91	0.6107
Metro - 2 Joinpoints	2	2008	2013	-5.4128*	-9.4 - 3.07	< 0.0001
Metro - 2 Joinpoints	3	2013	2020	2.0511*	0.66 - 4.54	0.0040
Non-Metro - 2 Joinpoints	1	1999	2014	-2.3813*	-3.63 - 1.12	0.0013
Non-Metro - 2 Joinpoints	2	2014	2018	14.3888	-1.77 - 33.21	0.0792
Non-Metro - 2 Joinpoints	3	2018	2020	-5.5088	-30.32 - 28.13	0.6959

Table 4. Joinpoint comparability analysis of temporal trends. *Comparison for American Indian or Alaska Native and Asian or Pacific Islander cannot be done as data was suppressed.

Model	Maximum Joinpoints (Kmax)	Test	Numerator DF	Denominator DF	Permutations	P-value	Interpretation
Sex-Combined Joinpoint model	2	Test of Parallelism	5	32	4,500	0.0100	Trends differ significantly between males and females
Urbanization-Combined Joinpoint model	2	Test of Parallelism	5	32	4,500	0.0036	Trends differ significantly between Metro and Non-Metro
Race-Combined Joinpoint model	2	Test of Parallelism	5	32	4,500	0.0002	Trends differ significantly between Black or African American and White

FIGURES

Figure 1. Overall age-adjusted mortality rates among adults aged 25+ in the United States, 1999–2020. *Indicates that the Annual Percentage Change (APC) is significantly different from zero at alpha=0.05 level.






Figure 2. Trends in sex-stratified age-adjusted Mortality rates among adults aged 25+ in the United States, 1999–2020. *Indicates that the Annual Percentage Change (APC) is significantly different from zero at alpha=0.05 level.




• Female - 2 Joinpoints	— 2007.0-2012.0 APC = -6.70*	▲ Male - 2 Joinpoints	— 2007.0-2015.0 APC = -2.82*
— 1999.0-2007.0 APC = -0.57	— 2012.0-2020.0 APC = 2.32*	— 1999.0-2007.0 APC = 0.47	— 2015.0-2020.0 APC = 4.96*

Figure 3. Trends in age-adjusted mortality rates stratified by race among adults aged 25+ years in the United States, 1999 to 2020. *Indicates that the Annual Percentage Change (APC) is significantly different from zero at alpha=0.05 level. Temporal trends for American Indian/Alaska Native and Asian or Pacific Islander are not displayed due to data suppression for counts <10, limiting reliable trend analysis.



• Black or African American - 2 Joinpoints	— 2014.0-2020.0 APC = 0.22	— 2008.0-2013.0 APC = -3.00
— 1999.0-2008.0 APC = -2.15	• White - 2 Joinpoints	— 2013.0-2020.0 APC = 3.34*
— 2008.0-2014.0 APC = -7.23*	— 1999.0-2008.0 APC = 0.05	

Figure 4. Temporal trends in age-adjusted mortality with intracerebral hemorrhage as the underlying cause of death and hypertension as a contributing cause, United States, 1999–2020.



■ Observed — 1999.0-2001.0 APC = 458.64* — 2001.0-2004.0 APC = -16.80* — 2004.0-2020.0 APC = -1.74*

* Indicates that the Annual Percent Change (APC) is significantly different from zero at the alpha = 0.05 level.
Final Selected Model: 1 Joinpoint.