



## OPEN ACCESS

## Letter to the editor

# Coronary artery systolic “milking” and “bridging” in Takotsubo syndrome: substrate or an epiphenomenon?

John E. Madias<sup>1,2,\*</sup>

<sup>1</sup>Icahn School of Medicine at Mount Sinai, New York, NY

<sup>2</sup>Division of Cardiology, Elmhurst Hospital Center, Elmhurst, NY

\*Email: madiasj@nychhc.org

To The Editor:

I enjoyed reading the case report by Spadotto et al.<sup>1</sup> published in issue 2 (2013) of the *Journal* about a 66-year-old woman with Takotsubo syndrome (TTS), involving the mid-apical segment of the left ventricle (LV), severe dynamic mid-ventricular gradient and mitral regurgitation, with systolic anterior motion of the anterior mitral leaflet caused by the hyperkinetic basal segments, with generation of a peak intra-ventricular gradient of 48 mmHg, and mild systolic “milking” of the mid to distal segment of the left anterior descending coronary artery (LAD). Cardiac magnetic resonance imaging (cMRI) showed intense myocardial edema of the mid-apical LV segments. The authors cite their previous report of “a recent clinically based study (which) showed that myocardial bridging of the LAD artery was detected in 76% of TTC patients either by angiography or MSCT”.<sup>2</sup> It is conceivable that this out of proportion high rate of “bridging”, and the occasionally reported systolic “milking” of the LAD in patients with TTS is due to an interplay of hypercontractility of the LV and the myocardial edema,<sup>3</sup> present in this condition. The issue can be easily resolved by coronary angiography (CA) (or MSCT), performed prior or after a bout of TTS in patients, who had/have a clinical indication for such a study, before or after their episode of TTS. In other words I am in doubt of the authors’ previous explanation that “myocardial bridging of the LAD is a frequent finding in TTS patients as revealed both by CA and, mostly, by MSCT, suggesting a role of myocardial bridging as potential substrate in the pathogenesis of TTS; probably this is an epiphenomenon of TTS state rather than an existing substrate prior to the inception of the TTS.

### Conflicts of Interest Disclosure

There is nothing to disclose.

### REFERENCES

- [1] Spadotto V, Elmaghawry M, Zorzi A, Migliore F, Marra MP. Apical ballooning with mid-ventricular obstruction: the many faces of Takotsubo cardiomyopathy. *Glob Cardiol Sci Pract*. 2013;2013(2):163–168, <http://dx.doi.org/10.5339/gcsp.2013.22>.
- [2] Migliore F, Maffei E, Perazzolo Marra M, Bilato C, Napodano M, Corbetti F, Zorzi A, Andres AL, Sarais C, Cacciavillani L, Favaretto E, Martini C, Seitun S, Cademartiri F, Corrado D, Iliceto S, Tarantini G. LAD coronary artery myocardial bridging and apical ballooning syndrome. *JACC Cardiovasc Imaging*. 2013;6(1):32–41, <http://dx.doi.org/10.1016/j.jcmg.2012.08.013>.
- [3] Perazzolo Marra M, Zorzi A, Corbetti F, De Lazzari M, Migliore F, Tona F, Tarantini G, Iliceto S, Corrado D. Apicobasal gradient of left ventricular myocardial edema underlies transient T-wave inversion and QT interval prolongation (Wellens’ ECG pattern) in Tako-Tsubo cardiomyopathy. *Heart Rhythm*. 2013;10(1):70–77, <http://dx.doi.org/10.1016/j.hrthm.2012.09.004>.

<http://dx.doi.org/10.5339/gcsp.2014.16>

Submitted: 4 April 2014  
Accepted: 9 April 2014  
© 2014 Madias, licensee  
Bloomsbury Qatar Foundation  
Journals. This is an open access  
article distributed under the terms  
of the Creative Commons  
Attribution license CC BY 4.0, which  
permits unrestricted use,  
distribution and reproduction in any  
medium, provided the original work  
is properly cited.