

Early detection of reduced left ventricular systolic function by 2D speckle tracking echocardiography in patients with primary mitral regurgitation in a Vietnamese cohort

Vu Anh Nguyen, Diem Thi Nguyen*, Linh Thi To Ong, An Viet Tran, Bao Lam Thai Tran, Chau Minh Tran

*ntdiem@ctump.edu.vn

DOI: <https://doi.org/10.21542/gcsp.2023.26>
VOLUME: 2023
ISSUE: 4
ARTICLE: 26

PMC PUBLICATION DATE: 30 Sept 2023
PMC COLLECTION DATE: 30 Sept 2023
PMC ARTICLE TYPE: research-article

ARTICLE TYPE: Research article
SUBMITTED: 20 MAY 2023
ACCEPTED: 12 AUG 2023
RUNNING HEAD: Nguyen et al. GCSP 2023:26

CITE THIS ARTICLE AS: Nguyen VA, Nguyen DT, Ong LTT, Tran AV, Tran BLT, Tran CM. Early detection of reduced left ventricular systolic function by 2D speckle tracking echocardiography in patients with primary mitral regurgitation in a Vietnamese cohort, Global Cardiology Science and Practice 2023:26 <https://doi.org/10.21542/gcsp.2023.26>

COPYRIGHT: 2023 The Author(s), licensee Magdi Yacoub Institute. This is an open access article distributed under the terms of the Creative Commons Attribution license CC BY-4.0, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Please note this is an 'Article in Press' and will be replaced with a typeset PDF & PubMed Central deposit in the coming weeks. Citation details and DOI will remain the same.

Early detection of reduced left ventricular systolic function by 2D speckle tracking echocardiography in patients with primary mitral regurgitation in a Vietnamese cohort

Vu Anh Nguyen¹, Diem Thi Nguyen^{2*}, Linh Thi To Ong³, An Viet Tran², Bao Lam Thai Tran², Chau Minh Tran²

1. Hue University of Medicine and Pharmacy, Hue, Viet Nam.
2. Can Tho University of Medicine and Pharmacy, Can Tho, Viet Nam.
3. Tam Duc Heart Hospital, Ho Chi Minh, Viet Nam.

[*ntdiem@ctump.edu.vn](mailto:ntdiem@ctump.edu.vn)

Abstract

Background: Mitral regurgitation (MR) is a common heart valve disease, causing many serious complications in several organ systems, especially the cardiovascular system. The 2D speckle tracking echocardiography (STE) is a new technique for detecting potential cardiac dysfunction when only tissue function abnormalities are present. The study aimed to assess left ventricular (LV) systolic function early by STE in patients with primary MR through global LV deformity along the global longitudinal strain (GLS).

Methods: An analytical cross-sectional study was performed on 46 patients with moderate to severe primary MR as recommended by the American Society of Echocardiography (ASE) 2017.

Results: The prevalence of patients with GLS reduction with ejection fraction (EF) >60%, New York Heart Association (NYHA) I, and left ventricular internal diameter systolic (LVIDs) <40mm was 38.1%, 35.7%, and 39.5%, respectively. 100% of patients with EF <60% and LVIDs ≥40mm had reduced GLS (<16%). The GLS index strongly correlates with the NYHA classification, degree of MR, EF, and echocardiographic parameters.

Conclusion: GLS index gives a significant sign in the early detection of cardiac function abnormalities before symptoms or other echocardiographic parameters in patients with MR.

Introduction

Mitral regurgitation (MR) is a valvular heart disease which has an increasing prevalence with age [1]. MR occurs in 10% of the general population [2]. In particular, left ventricular (LV) dysfunction will progress silently over many years with few or no symptoms. When the symptoms are clear, LV dysfunction may not be reversible, increasing morbidity and mortality rates despite surgery for mitral valve disease [3]. The key issue is determining when to intervene before the LV function reverses. Therefore, early detection of cardiac function abnormalities before symptoms can help clinicians have a better treatment strategy.

Finding reliable echocardiographic parameters for early detection of cardiac function abnormalities before symptoms or changes in classic echocardiographic parameters in patients with MR may help clinicians devise a better treatment strategy, deciding the appropriate timing of surgery to improve postoperative prognosis [3].

Two commonly used methods to evaluate LV function are M-mode echocardiography and Simpson's method. However, they only detect cardiac abnormalities when there is a complication of cardiac chamber dilation. Tissue Doppler ultrasonography can also help detect abnormalities of LV function early, but it is angle dependent. Many studies have shown that the echocardiographic markers of myocardial tissue detect potential cardiac dysfunction when there is only abnormal tissue function, no changes in cardiac morphology, and normal ejection fraction (EF). Simultaneously, the echocardiographic technique can assess cardiac function in different directions regardless of angle [4-6].

In Vietnam, studies using this technique for early detection of cardiac systolic function abnormalities in patients with MR, particularly those without surgical indications, are limited.

Materials and methods

Research objects

An analytical cross-sectional study of 46 patients with moderate to severe primary (organic) MR recommended by the American Society of Echocardiography (ASE) 2017 underwent cardiac Doppler echocardiography at Tam Duc Heart Hospital in Ho Chi Minh City from April 2019 to October 2020. The sample size is calculated with $n \geq 43$. Include $Z_{0.975} = 1.96$; $\alpha = 0.05$; $d = 0.09$; $p = 0.1$ [2].

Selection criteria: The patient was diagnosed with moderate to severe primary MR according to the recommendations of ASE 2017 by cardiac Doppler in Ho Chi Minh City [7]. The patient consented to participate in the study.

Exclusion criteria: Patients with secondary MR, MR due to myocardial infarction, MR combined with severe mitral stenosis and aortic valve stenosis, and heart failure with EF <40% (M-mode or Simpson). The patient had surgery to repair or replace the mitral valve. The patient has a bad ultrasound picture, and tachycardia (determined via STE frame rate). The patient did not consent to participate in the study.

Research content

Patient characteristics

General characteristics include (1) age; (2) gender (male and female); (3) causes of MR (mitral valve prolapse (MVP) and rheumatic heart disease).

Clinical characteristics of patients with MR include (1) degree of MR (severe and moderate were assessed by echocardiography and classified according to echocardiographic parameters: In patients diagnosed with mild MR, GLS has begun to decrease and from moderate to severe MR, GLS has already decreased) [7] (see Appendix A); (2) shortness of breath (including during exertion, during light exercise, and during paroxysmal or lying dyspnea); (3) New York Heart Association (NYHA) stands for classes of heart failure (NYHA I, NYHA II, NYHA III, NYHA IV) [7-9].

Cardiac Doppler echocardiography to classify primary and secondary MR is assessed through parameters including (1) ejection fraction (EF); (2) end-diastolic volume (EDV); (3) end-systolic volume (ESV); (4) left atrial volume (LAV); (5) pulmonary arterial pressures (PAPs).

Echocardiography parameters indicate that there are images of ischemic or non-ischemic cardiomyopathy with structural abnormalities or motion of the left ventricle, we classify the disease as secondary MR to exclude it.

According to the mitral regurgitation diagnostic criteria, MVP myxomatous changes, degenerative changes, infectious, inflammatory, and congenital causes are the main causes of primary MR. Based on the above causes, secondary mitral MR was excluded from the study [7] (Appendix B).

Characteristics of GLS index and correlation between GLS index and NYHA classification, degree of mitral regurgitation, EF, and echocardiographic parameters.

Characteristics of global longitudinal strain (GLS) index expressed through assessment of LV systolic function by echocardiography of 2D myocardium in patients with moderate to

severe primary MR assessed through GLS index through 3 longitudinal sections including (1) Four-chamber view at the apex (4CV); (2) Two-chamber view at the apex (2CV); (3) Three-chamber view at the apex of the heart (3CV).

Determining the correlations between NYHA classification, degree of MR, EF, and echocardiographic parameters with GLS include (1) Correlation between ejection fraction (EF) and GLS; (2) Correlation between the jet area/left atrium (LA) area (%) ratio with GLS; (3) Correlation between vena contracta (VC) and GLS; (4) Correlation between effective regurgitant orifice area (EROA) and GLS; (5) GLS index according to NYHA; (6) Correlation between end-diastolic volume (EDV) with GLS; (7) Correlation between end-systolic volume (ESV) and GLS; (8) Correlation between left atrial volume (LAV) and GLS; (9) Correlation between pulmonary arterial pressures (PAPs) and GLS.

Early assessment of LV systolic function by left ventricular GLS index in patients with primary MR

GLS decreased <16% with a cutoff value of -19.5% for GLS in the assessment of LV systolic function [6]. When GLS decreased <16%, EF >60%, NYHA I and LVIDs <40mm all had decreased expression. The prevalence of GLS reduction in patients with MR with NYHA I, EF >60%, and left ventricular internal diameter systolic (LVIDs) <40mm was determined through 3 parameters including (1) NYHA was divided into 4 classes NYHA I, NYHA II, NYHA III, NYHA IV; (2) LV EF includes 3 ranges of values >60%, 50-60%, <50%; (3) LVIDs include 2 ranges of values <40, ≥40 in patients with moderate and severe primary MR.

Data collection

Patients enrolled in the study underwent a data collection process including physical examination and questionnaire, routine M-mode echocardiography, Simpson, mitral flow Doppler, LAV, and deformity analysis of heart muscle.

The STE technique was performed by image acquisition with an average frame rate of 40- 80 fps [10]. To optimize imaging, the area to be examined should be placed at an intermediate depth, and the width of the area should be adjusted just enough. The axial view must pass through the apex or with the short axial view the left ventricular is round, the transverse deformation and torsion results are accurate [11]. Doppler ultrasound is measured in 4CV and 2CV at the apex, with the ultrasound beam aligned with the direction of motion of the area to be examined; preferably, the Doppler angle does not exceed 15°, with the most appropriate speed > 100 images/second and at least 4 QRS [11].

Data collection tools: The ultrasonic tissue marking process was performed using an Affiniti 70 ultrasound machine, with a frame rate of 1900 frame/s. QLAB 9.0 software, analyzed by aCMQ software. Probe S4-2, frequency 2-4 MHz.

Data analysis

Data analysis was performed using STATA14 software. Quantitative data are expressed as mean \pm SD and qualitative data as a percentage. Quantitative data were evaluated by an independent T-test, 1-way ANOVA (if >3 groups were compared) with a normal distribution while qualitative data were evaluated by a chi-squared test. Pearson's coefficient was used for the correlations. A p-value < 0.05 was considered to be statistically significant.

Ethics Committee Approval

The study was reported and approved by Hue University of Medicine and Pharmacy and the leadership of Tam Duc Heart Hospital, Ho Chi Minh City. (Decision No.1435 /QĐ-ĐHYD dated 30th July 2014).

Research diagram

The research was carried out according to the flow chart in Figure 1.




Figure 1. Study flow chart. EF, Ejection fraction; EDV, End-diastolic volume; ESV, End-systolic volume; GLS, Global longitudinal strain; LAV, Left atrial volume; NYHA, New York Heart Association; PAPs, Pulmonary arterial pressures.

Results

Table 1 describes the general characteristics and Doppler echocardiographic parameters of the study subjects. We found a predominance of women (56,5%) and an average age of 54.2 ± 14 years. The study recorded the indexes of EDV, ESV, EF%/M mode, EF%/Simpson (4CV), LAV, and PAPS with mean values of 100 ± 39 ml, 33 ± 17 ml, 69.6 ± 7.5 ml, respectively, 63.9 ± 6.1 ml, 65 ± 25 ml, and 33 ± 10 ml.

Table 1. General characteristics and Doppler echocardiographic parameters of the study subjects.

Characteristics	Patients (n=46) (\bar{X} + SD)
Gender	
Male	20 (43.5%)
Female	26 (56.5%)
Age (year)	54.2 ± 14.0
Doppler ultrasound parameters through the chest wall	
EDV (ml)	100 ± 39
ESV (ml)	33 ± 17
EF%/M mode	69.6 ± 7.5
EF%/Simpson (4CV)	63.9 ± 6.1
LAV (ml)	65 ± 25
PAPs (mmHg)	33 ± 10

EDV, End-diastolic volume; ESV, End-systolic volume; EF, Ejection fraction; 4CV, Four-chamber view; LAV, left atrial volume; PAPs, Pulmonary arterial pressures.

Table 2 describes the clinical characteristics and risk factors of the study subjects, revealing that 69.6% had severe MR and the remaining 30.4% had moderate MR. The cause of MR due to MVP accounts for 87.0%; the rest is rheumatic heart disease accounting for 13.0%. Subjects had the highest rate of dyspnea during exertion, 39.1%, dyspnea when mild exercise and paroxysmal dyspnea or lying down was 10.9% and 6.5%, respectively. Most of the study subjects had NYHA I heart failure class of 60.9%. The rate of heart failure in classes II, III, and IV, respectively, was 28.3%; 4.4%; and 6.5%.

Table 2. Clinical Characteristics and risk factors of study subjects

Characteristics	Patients (n=46)
Causes of MR	
MVP	40 (87.0%)
Rheumatic heart disease	6 (13.0%)
Total	46 (100%)
Degree of MR	
Severe	32 (69.6%)
Moderate	14 (30.4%)
Sudden onset of Dyspnea	
During exertion	18 (39.1%)
During light exercise	5 (10.9%)
During paroxysmal or lying dyspnea	3 (6.5%)
NYHA classification	

I	28 (60.9%)
II	13 (28.3%)
III	2 (4.4%)
IV	3 (6.5%)

MR, Mitral regurgitation; MVP, Mitral valve prolapse; NYHA, New York Heart Association.

Table 3 shows that there is a statistically significant difference ($p < 0.01$) in the mean GLS index in the severe MR group of $-15.4 \pm 2.2\%$ and the moderate MR group is $-19.7 \pm 2.0\%$. The echocardiography indicators showed a significant difference between the severe and moderate MR groups.

Table 3. Characteristics of ultrasonic parameters.

	Overall (n=46)	Degree of MR		p
		Severe (n=32)	Moderate (n=14)	
	$\bar{X} + SD$	$\bar{X} + SD$	$\bar{X} + SD$	
EF% /M - mode	69.6 ± 7.5	67.7 ± 7.6	74 ± 5.1	< 0.01
Simpson (4CV) EF%	63.9 ± 6.1	62.9 ± 6.3	66.4 ± 4.8	> 0.05
LAV (ml)	65 ± 25	75.1 ± 21.8	40.3 ± 14.6	<0.01
PAPs (mmHg)	33 ± 10	34.8 ± 10.8	28.9 ± 7.2	> 0.05
GLS (-)	16.7 ± 2.9	15.4 ± 2.2	19.7 ± 2.0	<0.01

EF, Ejection fraction; 4CV, Four-chamber view; GLS, Global longitudinal strain; LAV, Left atrial volume; MR, Mitral regurgitation; PAPs, Pulmonary arterial pressures.

Analysis of the results (Figure 2), shows a negative correlation between GLS and the jet area/LA area ratio ($r = -0.569$; $p < 0.01$), VC ($r = -0.592$; $p < 0.01$), and EROA ($r = -0.710$; $p < 0.01$). It demonstrates that as the jet area/LA area ratio, VC, and EROA increase, the GLS index decreases. The GLS index indicates that the more severe the MR, the lower the LV systolic function.




Figure 2 (A-C). Correlation between degree of mitral regurgitation and GLS. EROA, Effective regurgitant orifice area; GLS, Global longitudinal strain.

Figure 3 shows that there is a negative correlation between GLS with EDV ($r = -0.410$; $p < 0.01$), ESV ($r = -0.459$; $p < 0.01$), LAV ($r = -0.519$; $p < 0.01$), PAPs ($r = -0.342$, $p < 0.05$). Except that EF has a positive correlation with GLS ($r = 0.38$; $p < 0.05$), the absolute index of GLS increases when the EF index increases.




Figure 3 (A-E). Correlation between echocardiographic parameters with GLS. GLS, Global longitudinal strain.

The analysis of the relationship between the GLS index according to the NYHA classification (Figure 4) shows a statistically significant difference between the GLS and the NYHA class ($p < 0.05$). This means that the more severe the NYHA class, the lower the absolute index of GLS.




Figure 4. GLS index by NYHA. GLS, Global longitudinal strain; NYHA, New York Heart Association.

The results presented in Table 4 show that 35.7% of NYHA I patients had GLS reduction < 16%. Patients with NYHA II, III, and IV had a decrease in GLS <16% of 53.9%, 50.0%, and 66.7%, respectively. In patients with EF <50%, EF from 50-60%, 100% of patients with GLS decrease < 16%, and EF > 60% have 38.1% of patients with GLS decrease < 16%. In the group with LV dilatation with LVIDs ≥40mm, 100% of patients had a decrease in GLS <16%, and in the group with undilated left ventricles LVIDs <40mm, up to 39.5% of patients had a decrease in GLS <16%.

Table 4. Percentage of GLS reduction in patients with NYHA I, EF>60%, and LVIDs<40mm.

	Overall		GLS reduce <16*		Normal GLS ≥16	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
NYHA						
I	28	60.9	10	35.7	18	64.3
II	13	28.3	7	53.9	6	46.1
III	2	4.4	1	50.0	1	50.0
IV	3	6.5	2	66.7	1	33.3
EF%						
>60%	42	91.3	16	38.1	26	61.9
50-60%	3	6.5	3	100	0	0
<50%	1	2.2	1	100	0	0
LVIDs						
<40	43	93.5	17	39.5	26	60.5
≥40	3	6.5	3	100	0	0

EF, Ejection fraction; GLS, Global longitudinal strain; NYHA, New York Heart Association; LVIDs, Left ventricular internal diameter systolic.

* GLS decreased <16% with the cut-off value for GLS in assessing the LV systolic function of - 19.5%

[6].

Discussion

Patient characteristics

Our study's gender characteristics showed similar results to the studies of Reckefuss (male 49%, female 51%) and Nguyen Van Tha (male 47.6%, female 52.4%) [12, 13]. However, it differs from the Detaint (male 53.57%, female 46.43%), Grigioni (male 67%, female 33%), and Sutton (male 51.02%, female 48.92%) studies [13].

Our study's age characteristics are similar to those of other studies such as Rosenhek (55 ± 15), Singh (female 56 ± 13 , male 57 ± 11), Sutton (55 ± 19), data cross-sectional surveys, and other studies (median age 59 and 53 years) [13-16]. However, different from some studies such as Reckefuss (mean 42 years), Kobayashi, and Okura (mean 70.9 years) [12, 17].

Because of our study's smaller sample size compared to the research mentioned above, the age difference between men and women and the prevalence of the disease is not adequately reflected. Nguyen Van Tha's study differs from ours in terms of MR causes (93.7% MVP and 6.3% rheumatic heart disease) and MR degree (severe MR 50.8% and moderate MR 49.2%) and dyspnea (46% exertional dyspnea and 54% paroxysmal dyspnea at night or while lying down) [11]. Due to the characteristics of the Mekong Delta, people only go to the hospital when they have symptoms, as well as subjective symptoms of the patient.

Our NYHA classification of heart failure differs from some studies such as Nguyen Van Tha (no class I, 46% class II, 39.7% class III, 14.3% class IV), a study of over 150 patients who received mitral valve repair due to mitral valve insufficiency (60% class I, II and 40% class III, IV), and a prospective follow-up study on 551 patients had MR (27.3% class I, 38.7% class II, 34% class III) [18, 19]. Because most of the patients in our research sample, which was drawn from a clinic, were asymptomatic or only had minor heart failure symptoms, the rate of heart failure class I and II according to the NYHA is greater than it was for the authors' study.

Characteristics of GLS index and correlation between GLS index and NYHA classification, degree of mitral regurgitation, EF, and echocardiographic parameters

In our study, the severe MR group's GLS indexes tended to decrease statistically significantly (-15.4%) compared to the moderate regurgitation group (-19.7%) ($p < 0.001$), which showed similarities with the study of Ciro Santoro, Maurizio Galderisi, et al. (2019). In this study, the GLS index in the severe and mild MR group was -17.5% and -20.5%, respectively. ($p < 0.01$) [20]. Evaluation of the mean GLS index from our study reveals that it differs from other studies. The Norbert Reckefuss study gave an average GLS of -20.6% [10]. The average GLS in research on MR patients conducted by Ciro Santoro, Maurizio Galderisi, et al. (2019) was -19.8% [21]. The study of N. Bansal, A. Mercadante, et al. (2020) also gave an average GLS of -

22.5% [22]. Two other studies (2019) gave GLS indexes of -19.0% and -19.2%, respectively [23, 24]. Because normal GLS indexes differ depending on how the site is measured on the myocardium, the type of ultrasound machine used, and the software version used, published results vary significantly. Therefore, because the differences between vendors or software packages are still too great, most documents do not provide a uniformly normal reference value. In addition to software differences, GLS differences are also related to patients (age, gender, etc.) as well as dynamic factors (heart rate etc.)

Some guidelines approve GLS variation within the range of -20% considered normal in healthy individuals. According to Yingchoncharoen et al., LV GLS varied from -22.1% to -15.9%, with an average of -19.7%. Normal GLS according to Philips software and supplier (QLAB 7.1) is -18.9%. Several recent studies have reported lower normal values [25]. Most recently, according to the ACC (2018), in adults, a GLS index $< -16\%$ is considered abnormal, and a GLS index $> -18\%$ is normal [26].

Our research discovered an inverse correlation between GLS and VC, jet area/LA area ratio, and EROA, with correlation coefficients of -0.592, -0.569, and -0.710, respectively. According to many reports and literature, the higher the parameters such as VC, jet area/LA area ratio, and EROA, the more severe the MR, so the more severe the MR, the GLS index decreases. Our research found a negative correlation between GLS and VC ($r = -0.592$, $p = 0.01$). This finding is similar to that of Savas Dedeoglu et al., who discovered that in the moderate and severe mitral regurgitation groups, total left ventricular longitudinal strain correlated significantly with the vena contracta ($r = -0.858$; $p = 0.04$), and linear regression results revealed that the only independent association for LV longitudinal strain (model $r^2 = 0.311$) was vena contracta ($B = -0.858$; $p = 0.04$) [27]. Our research differs from Elena Kinova's study, which found a Pearson correlation between VCW and GLS ($r = 0.32$, $p = 0.015$) and found that the correlation was moderately positive right [28]. The difference might be explained by the two separate research populations.

The correlation between EF and GLS in our study is similar to that of other studies. Based on GLS, Y. Kobayashi; H. Okura, et al. (2017) discovered a link between EF and LV systolic function [17]. Research S.Yilmaze, R.G. Chelu, et al. (2020) discovered that EF correlates well with GLS [29]. Lima MSM, Villarraga HR, et al. (2017) also showed that the GLS of STE was strongly positively correlated with LVEF, particularly in LV systolic failure [30]. GLS was found to have a significant positive correlation with EF ($r = 0.33$; $p < 0.05$) in a study that used it to assess the diagnostic accuracy of LV systolic dysfunction by 2D tissue marker ultrasound to predict the severity of coronary artery disease [31]. In patients with preserved EF, reduced GLS ($< 16\%$) was associated with a 5.6-fold increased risk of death [32]. Overall, our findings are consistent with those of other authors.

Our study also discovered an inverse correlation between echocardiographic parameters (EDV, ESV, LAV, and PAPs) and GLS, with correlation coefficients of $r = -0.410$, -0.495 , -0.519 , and -0.342 , respectively. This finding is consistent with the findings of Ciro Santoro, Maurizio

Galderisi, et al. (2019) on subjects with MR and S. Yilmazer; R.G. Chelu, et al. (2020), who discovered that EDV and ESV were well correlated with GLS ($r > 0.61$, $p < 0.01$) [20, 29].

Our research was similar to other studies that found a statistically significant difference between GLS and NYHA classes ($p < 0.05$). Research by Mizuguchi, et al. also showed a decrease in GLS in hypertensive patients with LV hypertrophy in Asians; this decrease also increased with NYHA class with a statistically significant difference. In Kosmala's study, the GLS index gradually decreased from NYHA I to IV. The GLS index decreased in patients with heart failure with preserved EF compared to patients with hypertension but no heart failure in Nguyen Thi Diem's study, and this decline was even greater according to the NYHA class [6].

Early assessment of LV systolic function by left ventricular GLS index in patients with primary MR

Our study found that among NYHA I patients, 35.7% had a decrease in GLS $< 16\%$; even though the patient had no clinical symptoms, LV systolic function by GLS had already begun to decline. The NYHA classes II, III, and IV decreased the number of patients with GLS by 53.9%, 50.0%, and 66.7%, respectively. Through the study, we found that the prevalence of NYHA class from I to IV decreased, and the GLS decreased by $< 16\%$ (Especially NYHA I in the asymptomatic group, the GLS decreased early). This difference may suggest that GLS is statistically significant when it has already begun to decrease during the asymptomatic heart failure stage.

In our study, 100% of patients with EF reductions of $< 50\%$ and from 50%-60% had a GLS decrease of $< 16\%$. The findings revealed that in cases with EF at the transition stage of 50-60%, all subjects had reduced GLS. Although EF was only slightly reduced or normal, it was completely abnormal in GLS. In particular, in the group with EF $> 60\%$, up to 38.1% of patients had reduced GLS, corresponding to the compensated MR stage, but LV systolic function through the GLS index began to decrease. This demonstrates that GLS has a higher sensitivity than EF in detecting LV contractility abnormalities early.

According to Chin-in Lo et al., EF is insufficient to assess potential LV dysfunction in patients with preserved EF. In this case, GLS will superior EF in detecting early cardiac dysfunction in this group of patients [6, 33].

Ciro Santoro, Maurizio Galderisi, et al. (2019) found a correlation between GLS and EF and other echocardiographic parameters in 504 patients with mild to severe MR, which was similar to our findings. In particular, GLS was shown to be superior to EF in detecting LV systolic dysfunction in MR patients [20]. Our findings in normal patients show that even when EF has not yet decreased, GLS has already begun to decrease. This finding suggests that GLS has an early prognostic value in detecting LV dysfunction. To further demonstrate GLS's superiority, there is compelling evidence that LV-GLS has a higher predictive value than

LVEF in predicting LV dysfunction and serious cardiac events in many cardiovascular diseases [34].

Similarly, in the LV dilation group with LVIDs ≥ 40 mm, 100% of subjects had decreased GLS, while in the undilated left ventricle group LVIDs < 40 mm, 39.5% of subjects had also decreased GLS.

Limitations and implementations

Our study achieved many positive results, but it has some limitations. That is the sample size is small, and our research was conducted at only one center. To better understand the significance of this method in MR subjects, the sample size, and scope of the study must be expanded. More longitudinal studies with larger sample sizes or multicenter studies with longer follow-ups are required to assess the risk of heart failure between groups that have and do not have GLS changes on STE.

This study demonstrated that STE is superior to other ultrasound methods and that the GLS index on STE can be used to guide management, stratify prognosis, and determine an appropriate time for surgical intervention in patients with mitral regurgitation. Furthermore, in a large number of asymptomatic patients with severe primary mitral regurgitation and preserved ejection fraction, resting GLS may be shown to be a strong predictor of long-term prognosis.

However, assessing the group's decline in GLS on STE following surgical intervention is beyond the scope of the study. Therefore, further studies are needed to investigate the correlation of GLS decline on STE in the group before and after surgical intervention for primary mitral regurgitation.

Conclusions

Research results have shown that 2D STE helps in the early detection of abnormal manifestations of LV systolic function of the heart in patients with primary MR. In particular, for patients with no indication for surgery, the STE technique proved more effective than conventional ultrasound methods. The accuracy of diagnosing LV systolic dysfunction by 2D STE will increase further when combined with the LV GLS index.

Author statements

Conception: Vu A. Nguyen.

Design: Linh T. T. Ong.

Supervision: Diem T. Nguyen.

Funding: Diem T. Nguyen.

Materials: An V. Tran.

Data collection and/or processing: Linh T. T. Ong, Bao L. T. Tran, and Chau M. Tran.

Analysis and/or interpretation: An V. Tran.

Literature review: Diem T. Nguyen.

Writer: Bao L. T. Tran and Chau M. Tran.

Critical review: Vu A. Nguyen.

REFERENCES

- [1] Douedi S, Douedi H. Mitral Regurgitation. 2022 May 26. In: StatPearls [Internet]. *Treasure Island (FL): StatPearls Publishing*. 2022. PMID: 31985928.
- [2] Wu S, Chai A, Arimie S, Mehra A, Clavijo L, Matthews RV, Shavelle DM. Incidence and treatment of severe primary mitral regurgitation in contemporary clinical practice. *Cardiovasc Revasc Med*. 2018;19(8):960-963. doi:1016/j.carrev.2018.07.021.
- [3] Vietnam National Heart Association. 2008 Recommendations of the Vietnam Society of Cardiology on the diagnosis and treatment of valvular heart disease. *Journal of Vietnamese Cardiology*. 2008.
- [4] Dandel M, Hetzer R. Echocardiographic strain and strain rate imaging--clinical applications. *Int J Cardiol*. 2009;132(1):11-24. doi:1016/j.ijcard.2008.06.091.
- [5] de Simone G, Palmieri V. Diastolic dysfunction in arterial hypertension. *J Clin Hypertens (Greenwich)*. 2001;3(1):22-7. doi:1111/j.1524-6175.2001.00827.x.
- [6] Diem NT. Study of left ventricular function by speckle tracking echocardiography in patients with primary hypertension. *Specializing in Cardiology. Hue University of Medicine and Pharmacy*. 2017.
- [7] Zoghbi WA, Adams D, Bonow RO, Enriquez-Sarano M, Foster E, Grayburn PA, Hahn RT, Han Y, Hung J, Lang RM, Little SH, Shah DJ, Shernan S, Thavendiranathan P, Thomas JD, Weissman NJ. Recommendations for Noninvasive Evaluation of Native Valvular Regurgitation: A Report from the American Society of Echocardiography Developed in Collaboration with the Society for Cardiovascular Magnetic Resonance. *J Am Soc Echocardiogr*. 2017;30(4):303-371. doi:1016/j.echo.2017.01.007.
- [8] Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise J, Solomon S, Spencer KT, St John Sutton M, Stewart W. Recommendations for chamber quantification. *Eur J Echocardiogr*. 2006;7(2):79-108. doi:1016/j.euje.2005.12.014.
- [9] Vinh PN. Cardiovascular pathology, volume II. *Specializing in Cardiology. Medical Publishing House*. 2008.
- [10] Marwick TH, Leano RL, Brown J, Sun JP, Hoffmann R, Lysyansky P, Becker M, Thomas JD. Myocardial strain measurement with 2-dimensional speckle-tracking echocardiography: definition of normal range. *JACC Cardiovasc Imaging*. 2009;2(1):80-4. doi: 10.1016/j.jcmg.2007.12.007.
- [11] Mor-Avi V, Lang RM, Badano LP, Belohlavek M, Cardim NM, Derumeaux G, Galderisi M, Marwick T, Nagueh SF, Sengupta PP, Sicari R, Smiseth OA, Smulevitz B, Takeuchi M, Thomas JD, Vannan M, Voigt JU, Zamorano JL. Current and evolving echocardiographic techniques for

- the quantitative evaluation of cardiac mechanics: ASE/EAE consensus statement on methodology and indications endorsed by the Japanese Society of Echocardiography. *J Am Soc Echocardiogr*. 2011;24(3):277-313. doi: 10.1016/j.echo.2011.01.015.
- [12] Reckefuss N, Butz T, Horstkotte D, Faber L. Evaluation of longitudinal and radial left ventricular function by two-dimensional speckle-tracking echocardiography in a large cohort of normal probands. *Int J Cardiovasc Imaging*. 2011;27(4):515-26. doi:1007/s10554-010-9716-y.
- [13] Tha NV. Study on NT-pro BNP concentration in patients with heart failure due to mitral valve regurgitation. *Hue University of Medicine and Pharmacy*. 2017.
- [14] Di Stefano C, Bruno G, Arciniegas Calle MC, Acharya GA, Fussner LM, Ungprasert P, Cooper LT, Blauwet LA, Ryu JH, Pellikka PA, Carmona Porquera EM, Villarraga HR. Diagnostic and predictive value of speckle tracking echocardiography in cardiac sarcoidosis. *BMC cardiovascular disorders*. 2020;20:21. doi:1186/s12872-019-01323-0.
- [15] Kleijn SA, Aly MF, Terwee CB, van Rossum AC, Kamp O. Reliability of left ventricular volumes and function measurements using three-dimensional speckle tracking echocardiography. *Eur Heart J Cardiovasc Imaging*. 2012;13(2):159-68. doi:1093/ejehocardi/jer174.
- [16] van der Bijl P, Bootsma M, Hiemstra YL, Ajmore Marsan N, Bax JJ, Delgado V. Left ventricular 2D speckle tracking echocardiography for detection of systolic dysfunction in genetic, dilated cardiomyopathies. *Eur Heart J Cardiovasc Imaging*. 2019;20(6):694-699. doi:1093/ehjci/jey169.
- [17] Kobayashi Y, Okura H, Kobayashi Y, Fukuda S, Hirohata A, Yoshida K. Left ventricular myocardial function assessed by three-dimensional speckle tracking echocardiography in Takotsubo cardiomyopathy. *Echocardiography*. 2017;34(4):523-529. doi:1111/echo.13492.
- [18] Abid A, Denguir R, Hakim M, Gharsallah N, Ghedira F, Meddeb I, Kalfat T, Khayati A. [Relevant factors in mitral valve reconstructive surgery. Report of 150 cases]. *La Tunisie medicale*. 2002;80(4):183-187. PMID: 12416352
- [19] Carpentier A, Chauvaud S, Fabiani JN, Deloche A, Relland J, Lessana A, D'Allaines C, Blondeau P, Piwnica A, Dubost C. Reconstructive surgery of mitral valve incompetence: ten-year appraisal. *J Thorac Cardiovasc Surg*. 1980;79(3):338-48. doi:1016/S0022-5223(19)37942-5.
- [20] Santoro C, Galderisi M, Esposito R, Buonauro A, Monteagudo JM, Sorrentino R, Lembo M, Fernandez-Golfín C, Trimarco B, Zamorano JL. Global longitudinal strain is a hallmark of cardiac damage in mitral regurgitation: the Italian arm of the European Registry of mitral regurgitation (EuMiClip). *Cardiovasc Ultrasound*. 2019;17(1):28. doi:1186/s12947-019-0178-7.
- [21] Vu NA. Echocardiography - diagnostic update. *Hue University Publishing House*. 2018;192-237.
- [22] Bansal N, Mercadante A, Rochelson E, Mahgerefteh J, Clark BC. Speckle Tracking Echocardiography in Pediatric Patients with Premature Ventricular Contractions. *Pediatr Cardiol*. 2020;41(8):1587-1593. doi:1007/s00246-020-02415-x.
- [23] Rong LQ, Yum B, Abouzeid C, Palumbo MC, Brouwer LR, Devereux RB, Girardi LN, Weinsaft JW, Gaudino M, Kim J. Echocardiographic predictors of intraoperative right ventricular dysfunction: a 2D and speckle tracking echocardiography study. *Cardiovascular ultrasound*. 2019;17:11. doi:1186/s12947-019-0161-3.
- [24] Zamirian M, Afsharizadeh F, Moaref A, Abtahi F, Amirmoezi F, Attar A. Reduced myocardial reserve in cirrhotic patients: an evaluation by dobutamine stress speckle tracking and tissue Doppler imaging (TDI) echocardiography. *J Cardiovasc Thorac Res*. 2019;11(2):127-131. doi:15171/jcvtr.2019.22.

- [25] Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, Flachskampf FA, Foster E, Goldstein SA, Kuznetsova T, Lancellotti P, Muraru D, Picard MH, Rietzschel ER, Rudski L, Spencer KT, Tsang W, Voigt JU. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr.* 2015;28(1):1-39.e14. doi:10.116/j.echo.2014.10.003
- [26] Yang H, Wright L, Negishi T, Negishi K, Liu J, Marwick Thomas H. Research to Practice. *JACC: Cardiovascular Imaging.* 2018;11(8):1196-1201. doi:10.116/j.jcmg.2018.07.005.
- [27] Dedeoğlu S, Bornaun H, dedeoğlu R. The use of left atrial strain to predict left ventricular functions in asymptomatic children with mitral valve regurgitation. 2023. doi:10.22541/au.168233067.75068451/v1.
- [28] Kinova E, Spasova N, Borizanova A, Goudev A. Torsion Mechanics as an Indicator of More Advanced Left Ventricular Systolic Dysfunction in Secondary Mitral Regurgitation in Patients with Dilated Cardiomyopathy: A 2D Speckle-Tracking Analysis. *Cardiology.* 2018;139(3):187-196. doi: 10.1159/000485967.
- [29] van den Hoven AT, Yilmazer S, Chelu RG, van Grootel RWJ, Minderhoud SCS, Bous LR, van Berendoncks AM, Duijinhouwer AL, Siebelink HMJ, van den Bosch AE, Budde RPJ, Roos-Hesselink JW, Hirsch A. Left ventricular global longitudinal strain in bicuspid aortic valve patients: head-to-head comparison between computed tomography, 4D flow cardiovascular magnetic resonance and speckle-tracking echocardiography. *International Journal of Cardiovascular Imaging.* 2020;36(9):1771 - 1780. doi:10.1007/s10554-020-01883-9.
- [30] Lima MSM, Villarraga HR, Abduch MCD, Lima MF, Cruz Cbbv, Sbano JCN, Voos MC, Mathias W Junior, Tsutsui JM. Global Longitudinal Strain or Left Ventricular Twist and Torsion? Which Correlates Best with Ejection Fraction?. *Arq Bras Cardiol.* 2017;109(1)23-29. doi:10.5935/abc.20170085.
- [31] Radwan H, Hussein E. Value of global longitudinal strain by two dimensional speckle tracking echocardiography in predicting coronary artery disease severity. *Egypt Heart J.* 2017;69(2):95-101. doi:10.116/j.ehj.2016.08.001.
- [32] Krishnasamy R, Isbel NM, Hawley CM, Pascoe EM, Burrage M, Leano R, Haluska BA, Marwick TH, Staton T. Left Ventricular Global Longitudinal Strain (GLS) Is a Superior Predictor of All-Cause and Cardiovascular Mortality When Compared to Ejection Fraction in Advanced Chronic Kidney Disease. *PLoS One.* 2015;10(5):e0127044. doi:1371/journal.pone.0127044.
- [33] Cam VN. Investigate the effect of blood pressure on left ventricular function by longitudinal muscle tension. *Pham Ngoc Thach Medical University.* 2019.
- [34] Kalam K, Otahal P, Marwick TH. Prognostic implications of global LV dysfunction: a systematic review and meta-analysis of global longitudinal strain and ejection fraction. *HEART.* 2014;100(21):1673-1680. doi:10.1136/heartjnl-2014-305538.